

Section 4.2


Maximum and Minimum Values

- (1) Absolute and Local Extrema
- (2) The Extreme Value Theorem
- (3) The Closed Interval Method

Local Extrema


A function f has a **local maximum** at c if $f(c) \geq f(x)$ for x "near" c .
That is, $f(c) \geq f(x)$ for all x in some open interval containing c .

A function f has a **local minimum** at c if $f(c) \leq f(x)$ for x "near" c .




Local Extrema

- The term “extremum” is shorthand for “maximum or minimum.”
- If f has a local extremum at c , then $y = f(c)$ is a **local extreme value** and $(c, f(c))$ is a **local extreme point**.
- An endpoint of the domain f of cannot be a local extremum, because it cannot be contained in any open interval in the domain.
- A function does not necessarily have to have any local extrema:




No maximum.

No minimum.



No maximum.

Minimum value is 0.



No minimum.


Maximum value is 2.

Absolute Extrema

A function f has an **absolute maximum** at c if $f(c) \geq f(x)$ for **all** x .

A function f has an **absolute minimum** at c if $f(c) \leq f(x)$ for **all** x .

- Unless it is an endpoint, each absolute extremum is also a local extremum.




Example 1: Absolute Extrema

A function can have **at most one** absolute maximum value, but **any** number of absolute maximum points.


Example 1(a): $f(x) = x^2$ has **no absolute maximum**, because x^2 can be arbitrarily large.

Example 1(b): $f(x) = -x^2$ has **absolute maximum** value 0 at the point $(0,0)$.

Example 1(c): $f(x) = \cos(x)$ has absolute maximum value 1 and infinitely many absolute maximum points: $(k\pi, 1)$ where k is any even integer.



Example 2: Local and Absolute Maxima and Minima





Critical Numbers and Fermat's Theorem

A number c in the domain of f is called a **critical number** if either $f'(c) = 0$ or $f'(c)$ does not exist.

Fermat's Theorem

If f has a local extremum at $x = c$, and $f'(c)$ exists, then $f'(c) = 0$.

That is, if f has a local max or min at c , then c is a critical number of f .






Critical Numbers

Fermat's Theorem

If f has a local extremum at $x = c$, and $f'(c)$ exists, then $f'(c) = 0$.







That is, if f has a local max or min at c , then c is a critical number of f .

On the other hand, not every critical point must be a local max or min.



The Extreme Value Theorem

If f is continuous on a **closed** interval $[a, b]$, then f attains an absolute maximum value $f(c)$ and an absolute minimum value $f(d)$ at some numbers c and d in $[a, b]$.



(A) A discontinuous function on interval $[a, b]$ has a min but no max on interval $[a, b]$.

(B) A continuous function on interval (a, b) has no min or max on open interval (a, b) .

(C) Every continuous function on a closed interval $[a, b]$ has both a min and a max on $[a, b]$.

The Closed Interval Method

The Extreme Value Theorem

If f is continuous on a **closed** interval $[a, b]$, then f attains an absolute maximum value $f(c)$ and an absolute minimum value $f(d)$ at some numbers c and d in $[a, b]$.

How do we systematically find the absolute extrema?

The Closed Interval Method

To find the absolute extreme values of a continuous function f on a closed interval $[a, b]$:

- (1) Find the values of f at the **critical numbers** in (a, b) .
- (2) Find the values of f at the **endpoints** (namely a and b).
- (3) Compare the y -values. The largest value is the absolute maximum value; the smallest value is the absolute minimum value.

The Closed Interval Method

Example 3: Find the absolute extrema of $f(x) = 2x^3 - 15x^2 + 24x + 7$ on the closed interval $[0,6]$.

Note that f is a polynomial, so it is continuous everywhere.



Finding Extrema

Example 4: Find the absolute extrema of $h(x) = x^{4/5}(x-4)^2$ on $[1, 5]$.



Finding Extrema

Example 4: Find the absolute extrema of $h(x) = x^{4/5}(x-4)^2$ on $[1, 5]$.

