

## Section 4.3

### The Mean Value Theorem


- (1) Rolle's Theorem and the Mean Value Theorem
- (2) The First Derivative Test

## Rolle's Theorem

Suppose that  $f$  is a function such that

- (I)  $f$  is continuous on  $[a, b]$ ,
- (II)  $f$  is differentiable on  $(a, b)$ ,
- (III)  $f(a) = f(b)$ .

Then there exists a value  $c$  in  $(a, b)$  such that  $f'(c) = 0$ .



Equivalently: there is a value  $c$  in  $(a, b)$  such that the **tangent line** at  $(c, f(c))$  is **parallel** to the **secant line** from  $(a, f(a))$  to  $(b, f(b))$ .

## Rolle's Theorem

Suppose that  $f$  is a function such that

- (I)  $f$  is continuous on  $[a, b]$ ,
- (II)  $f$  is differentiable on  $(a, b)$ ,
- (III)  $f(a) = f(b)$ .

Then there exists a value  $c$  in  $(a, b)$  such that  $f'(c) = 0$ .

Equivalently: there is a value  $c$  in  $(a, b)$  such that the **tangent line** at  $(c, f(c))$  is **parallel** to the **secant line** from  $(a, f(a))$  to  $(b, f(b))$ .



What if we drop the assumption that  $f(a) = f(b)$ ?

## The Mean Value Theorem (MVT)

Suppose that  $f$  is a function that is **(I)** continuous on  $[a, b]$ , and **(II)** differentiable on  $(a, b)$ .

Then there exists a value  $c$  in  $(a, b)$  such that  $f'(c) = \frac{f(b) - f(a)}{b - a}$ .

**Idea:** Take the secant line from  $(a, f(a))$  to  $(b, f(b))$  and slide it until it becomes a tangent line.



# Mean Value Theorem: Examples

**Example 1:** Let  $f(x) = x^3 - x$  on the interval  $[0, 2]$ .



# Mean Value Theorem: Examples

**Example 2:** Let  $f(x) = x^4 - x^2 - 3x$  on the interval  $[0, 2]$ .



# Mean Value Theorem: Examples

**Example 3:** Let  $f(x) = \tan(x)$ .



# Mean Value Theorem: Examples

**Example 4:** Find  $M$  and  $m$  such that  $m \leq f(-2) \leq M$  if  $f$  is a function where  $f(1) = 3$  and  $-1 \leq f'(x) \leq 4$  for all  $x$ .





## Consequences of the Mean Value Theorem

**Theorem 1:** If  $f'(x) = 0$  for all  $x$  in  $(a, b)$ , then  $f$  is constant on  $(a, b)$ .

**Theorem 2:** If  $f'(x) > 0$  for all  $x$  in  $(a, b)$ , then  $f$  is increasing on  $(a, b)$ .

**Theorem 3:** If  $f'(x) = g'(x)$  for all  $x$  in  $(a, b)$ , then  $f(x) = g(x) + C$ , where  $C$  is some constant.

**Proof of Theorem 1:** Suppose that  $A, B$  are arbitrary numbers with  $a < A < B < b$ .

Since  $f'$  exists for all values in  $(a, b)$ ,  $f$  is continuous and differentiable on  $(a, b)$  and MVT holds on the interval  $[A, B]$ .

Therefore, there exists  $c$  in  $[A, B]$  such that

$$f'(c) = \frac{f(B) - f(A)}{B - A}$$

But  $f'(c) = 0$ , so  $f(B) - f(A) = (B - A)0 = 0$ .

## Consequences of the Mean Value Theorem

**Theorem 1:** If  $f'(x) = 0$  for all  $x$  in  $(a, b)$ , then  $f$  is constant on  $(a, b)$ .

**Theorem 2:** If  $f'(x) > 0$  for all  $x$  in  $(a, b)$ , then  $f$  is increasing on  $(a, b)$ .

**Theorem 3:** If  $f'(x) = g'(x)$  for all  $x$  in  $(a, b)$ , then  $f(x) = g(x) + C$ , where  $C$  is some constant.

**Proof of Theorem 2:** We need to prove that if  $a < A < B < b$ , then  $f(A) < f(B)$ . Again, apply the MVT to the interval  $[A, B]$ . The conclusion is that there exists  $c$  in  $[A, B]$  such that

$$f'(c) = \frac{f(B) - f(A)}{B - A}.$$

But  $f'(c) > 0$  and  $B - A > 0$ , so  $f(B) - f(A) > 0$ , i.e.,  $f(B) > f(A)$ .

## Consequences of the Mean Value Theorem

**Theorem 1:** If  $f'(x) = 0$  for all  $x$  in  $(a, b)$ , then  $f$  is constant on  $(a, b)$ .

**Theorem 2:** If  $f'(x) > 0$  for all  $x$  in  $(a, b)$ , then  $f$  is increasing on  $(a, b)$ .

**Theorem 3:** If  $f'(x) = g'(x)$  for all  $x$  in  $(a, b)$ , then  $f(x) = g(x) + C$ , where  $C$  is some constant.

**Proof of Theorem 3:** Let  $h(x) = f(x) - g(x)$ .

Since  $f'(x) = g'(x)$ , it follows that  $h'(x) = 0$  on  $(a, b)$ .

Now Theorem 1 implies that  $h(x) = C$  on  $(a, b)$ , where  $C$  is some constant.


Therefore  $f(x) = g(x) + C$ .

# The First Derivative Test

## First Derivative Test for Local Extrema

Suppose that  $c$  is a critical number of a continuous function  $f$ .

- (I) If  $f'$  changes from positive to negative at  $c$ , then  $f$  has a local maximum at  $c$ .
- (II) If  $f'$  changes from negative to positive at  $c$ , then  $f$  has a local minimum at  $c$ .
- (III) If  $f'$  does not change sign at  $c$ , then  $c$  is not a local extremum.



**Example 5:** Find the local extrema of  $f$  if  $f'(x) = (x-4)^3(x+3)^7(x-2)^6$ .



**Example 6:** Find the local extrema of the function  $f(x) = \frac{(x+1)^2}{x(x-2)}$ .

