

# Section 5.1

## Areas and Distances

- (1) The Area Problem
- (2) The Distance Problem
- (3) Summation Notation

# Area

Area is a measure of the size of 2-dimensional shapes.

Area is preserved under cutting, gluing, sliding, and rotating.

There are standard formulas for the areas of common shapes:

**Rectangle:**  $A = bh$

**Triangle:**  $A = \frac{1}{2}bh$

**Circle:**  $A = \pi r^2$

But what about more complicated shapes?

# The Area Problem

The motivation for this chapter is the problem of calculating the area of more general regions, such as the area under the graph of a function  $y = f(x)$ .


When we studied tangent lines, we soon discovered that we needed to use **limits** to calculate them in a mathematically rigorous way. This led to the concept of a **derivative**.

Similarly, calculating area in a rigorous way will also require **limits** and will lead us to a new mathematical concept: the **integral**.

# The Area Problem

Let  $f(x)$  be continuous and positive on a closed interval  $[a, b]$ .

**What is the area of the region bounded by the graph of  $f(x)$ , the vertical lines  $x = a$  and  $x = b$ , and the  $x$ -axis?**



# The Area Problem


The area  $A$  under the graph of  $f$  between  $x = a$  and  $x = b$  can be **approximated** as the total area of  $n$  rectangles.

- Divide the domain  $[a, b]$  into  $n$  segments of length  $\Delta x = \frac{b-a}{n}$ .
- Inside each segment, choose a value  $x_i$ .
- Form a rectangle of height  $f(x_i)$  on each segment.

Then  $A \approx f(x_1) \Delta x + f(x_2) \Delta x + \dots + f(x_n) \Delta x.$

by Joseph Phillip Brennan  
Jila Niknejad

**Example 1:** Approximate the area under  $y = x^2$  on  $[1,4]$  using 6 segments.



## Area Expressed as a Limit

The area  $A$  under the graph of  $f$  between  $x = a$  and  $x = b$  can be **approximated** as the total area of  $n$  rectangles:

$$A \approx \overbrace{f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_n)\Delta x}^{R_n}$$

As  $n$  gets larger and larger, the approximation  $R_n$  gets better and better.

The **exact** area is given by a **limit**.

The **area**  $A$  under the graph of a continuous function  $f$  between  $x = a$  and  $x = b$  equals the limit of the sum of the areas of approximating rectangles:

$$A = \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} (f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_n)\Delta x)$$

# Calculating Distance

Let  $v(t)$  be the velocity of an object at time  $t$ .


The area under the graph of  $v(t)$  on a time interval  $[a, b]$  measures the **net distance traveled**, or **displacement**, between times  $a$  and  $b$ .

Note that the units:


$$\begin{aligned}\text{Units of area under the graph of } v(t) &= \text{units of } t \times \text{units of } v(t) \\ &= \text{time} \times \frac{\text{distance}}{\text{time}} \\ &= \text{distance}.\end{aligned}$$



**Example 2(a):** If  $v = v_0$  on  $[a, b]$ , then the region under the graph is a rectangle with area  $v_0(b - a)$ .



**Example 2(b):** An object starts at rest and accelerates at a constant rate of  $1 \text{ m/s}^2$  for 10 seconds. Then  $v(t) = t \text{ m/s}$ .  
Displacement = area under the curve =  $\frac{1}{2}(10 \text{ s})(10 \text{ m/s}) = 50 \text{ m}$ .



**Example 3:** You are driving across Missouri. In order to stay awake, you estimate how far you have traveled from your speedometer readings:

2:00 PM	70 mph (the speed limit)
2:15 PM	65 mph (up a small hill)
2:30 PM	75 mph (down the hill)
2:45 PM	55 mph (careful, is that a speed trap?)
3:00 PM	80 mph (vroom!)

You can now estimate<sup>1</sup> the maximum and minimum possible distance you have traveled during this hour:

$$\text{Max: } \frac{1}{4}(70) + \frac{1}{4}(75) + \frac{1}{4}(75) + \frac{1}{4}(80) = 75 \text{ miles}$$

$$\text{Min: } \frac{1}{4}(65) + \frac{1}{4}(65) + \frac{1}{4}(55) + \frac{1}{4}(55) = 60 \text{ miles}$$

The actual distance traveled is somewhere between these two estimates.

---

<sup>1</sup>Assuming that in each 15-minute interval, your max and min speeds occur at the endpoints.

# Summation Notation

We have encountered expressions like

$$f(x_1)\Delta x + f(x_2)\Delta x + \dots + f(x_n)\Delta x$$

that are **sums** of many similar-looking terms. We need a notation for writing sums in a compact form.

## Summation Notation

The notation  $\sum_{j=m}^n a_j$  means  $a_m + a_{m+1} + a_{m+2} + \dots + a_{n-1} + a_n$ .

- $\Sigma$  is the Greek letter Sigma (mnemonic for “sum.”)
- The notation  $\sum_{j=m}^n$  tells us to start at  $j = m$  and to end at  $j = n$ .
- $a_j$  is called the **general term** and  $j$  is the **summation index**.

## Summation Notation

The notation  $\sum_{j=m}^n a_j$  means  $a_m + a_{m+1} + a_{m+2} + \dots + a_{n-1} + a_n$ .

Examples:

$$\sum_{j=1}^{100} j = 1 + 2 + 3 + \dots + 100$$

$$\sum_{j=4}^{785} j^2 = 4^2 + 5^2 + \dots + 785^2$$

$$\sum_{j=4}^6 (j^3 - j - 1) = (4^3 - 4 - 1) + (5^3 - 5 - 1) + (6^3 - 6 - 1)$$

# Summation Notation and Area

## Summation Notation

The notation  $\sum_{j=m}^n a_j$  means  $a_m + a_{m+1} + a_{m+2} + \dots + a_{n-1} + a_n$ .

Therefore, our **estimate** for the area under the graph of a continuous, positive function  $f(x)$  on an interval  $[a, b]$  is

$$R_n = f(x_1)\Delta x + \dots + f(x_n)\Delta x = \sum_{j=1}^n f(x_j)\Delta x$$

and the **exact area** is

$$\begin{aligned} A &= \lim_{n \rightarrow \infty} R_n = \lim_{n \rightarrow \infty} (f(x_1)\Delta x + \dots + f(x_n)\Delta x) \\ &= \lim_{n \rightarrow \infty} \sum_{j=1}^n f(x_j)\Delta x. \end{aligned}$$

# Properties of Summations

If you understand addition, you understand summation!

$$\sum_{j=m}^n (a_j \pm b_j) = \left( \sum_{j=m}^n a_j \right) \pm \left( \sum_{j=m}^n b_j \right)$$

$$\sum_{j=m}^n (ca_j) = c \sum_{j=m}^n a_j \quad (\text{for any constant } c)$$

$$\sum_{j=m}^n c = c(n - m + 1)$$

# Properties of Summations

For example:

$$\begin{aligned}\sum_{j=1}^{1000} (3j^2 - 5j + 3) &= \left( \sum_{j=1}^{1000} 3j^2 \right) - \left( \sum_{j=1}^{1000} 5j \right) + \left( \sum_{j=1}^{1000} 3 \right) \\ &= 3 \left( \sum_{j=1}^{1000} j^2 \right) - 5 \left( \sum_{j=1}^{1000} j \right) + 3000\end{aligned}$$

Fortunately, there are nice formulas for the sum of the first  $n$  numbers, squares, cubes, fourth powers, ...

# Summation Formulas

- $\sum_{j=1}^n j = 1 + 2 + \dots + (n-1) + n = \frac{n(n+1)}{2}$
- $\sum_{j=1}^n j^2 = 1^2 + 2^2 + \dots + (n-1)^2 + n^2 = \frac{n(n+1)(2n+1)}{6}$
- $\sum_{j=1}^n j^3 = 1^3 + 2^3 + \dots + (n-1)^3 + n^3 = \frac{n^2(n+1)^2}{4}$

You don't have to memorize these formulas, but the first one has a very elegant explanation!




# The Sum of the First $N$ Integers



$$1 + 2 + \dots + N = \frac{N \times (N + 1)}{2}$$

# The Sum of the Cubes of the First N Integers



$$1^3 + 2^3 + \dots + N^3 = (1 + 2 + \dots + N)^2$$